



TITLE:

A Note on 2-Fold Branched Covers of S^3 (低次元多様体の構造と分類について)

AUTHOR(S):

KANENOBU, TAIZO

CITATION:

KANENOBU, TAIZO. A Note on 2-Fold Branched Covers of S^3 (低次元多様体の構造と分類について). 数理解析研究所講究録 1981, 417: 28-35

ISSUE DATE:

1981-02

URL:

<http://hdl.handle.net/2433/102495>

RIGHT:

A Note on 2-fold Branched Covers of S^3

Taizo KANENOBU

Kobe University

1. Introduction

If a knot $K \subset S^3$ is amphicheiral, then the 2-fold covering space branched over K is symmetric (has an orientation reversing diffeomorphism). In this note we show that the converse is not true; it solves Problem 1.23 of [K], which is suggested by Montesinos [M1].

Theorem. There exist non-amphicheiral knots whose 2-fold branched covering spaces are symmetric.

In Section 2, we will construct a composite knot satisfying the properties of Theorem, and in Section 3, a prime knot.

2. A Composite Knot


Let K_1 and K_2 be two knots. If there exists an orientation preserving homeomorphism of S^3 which maps K_1 onto K_2 , then we will write $K_1 \approx K_2$.

Lemma 1. Let K_1 and K_2 be two prime knots such that $K_1 \approx r(K_2)$ where r is an orientation reversing homeomorphism of S^3 . Then the

composite knot $K_1 \# K_2$ is amphicheiral if and only if both K_1 and K_2 are amphicheiral.

Proof. If $K_1 \# K_2$ is amphicheiral, then $K_1 \# K_2 \approx r(K_1 \# K_2) = r(K_1) \# r(K_2)$. It follows from the unique decomposition theorem [Sc] and $K_1 \not\approx r(K_2)$ that $K_1 \approx r(K_1)$ and $K_2 \approx r(K_2)$. The converse is clear.

Let K_1 be the pretzel knot (see [Tr] or [P]) of type $(p_1, \dots, p_i, \dots, p_j, \dots, p_n)$, denoted by $K(p_1, \dots, p_i, \dots, p_j, \dots, p_n)$, where n (≥ 5) is an odd integer and p_1, p_2, \dots, p_n (≥ 3) are distinct and odd integers. Let $K_2 = K(p_1, \dots, p_j, \dots, p_i, \dots, p_n)$, where $i \neq j$. By the theorem in §2 of [M1], the 2-fold covering spaces of S^3 branched over K_1 and K_2 are the same Seifert fiber space $(0 \cup 0 \mid 0; (p_1, 1), (p_2, 1), \dots, (p_n, 1))$ [Se], or the manifold with the following surgery presentation (see [R, Chapter 9]):



This manifold is prime by Theorem 7.1 and Lemma 10.2 of [W1]. Hence by the main result of [W2], K_1 and K_2 are prime knots. By the classification theorem of pretzel knots [P, p.56], we have $K_1 \not\approx K_2$. Since the signatures of K_1 and K_2 are both $n-1$ ($\neq 0$) [P, p.71], K_1 and K_2 are non-amphicheiral [R, p.217]. It follows from Lemma 1 that $K_1 \# r(K_2)$ is non-amphicheiral.

On the other hand, the 2-fold covering space of S^3 branched over $K_1 \# r(K_2)$ is clearly symmetric. Therefore $K_1 \# r(K_2)$ is the desired composite knot.

Remark. We can also construct such a composite knot by using the examples of prime knots in [BGM] and [Ta] instead of pretzel knots.

3. A Prime Knot

Let M be the manifold which is obtained by Dehn surgery in the link L according to the diagram of Fig. 1. Since the figure-eight knot, which is the sublink of L , is amphicheiral, it is easily verified that M is symmetric. The link L is strongly-invertible because the symmetry with respect to the axis E leaves L invariant. Using the method of [M2], it can be shown that M is a 2-fold covering space branched over the knot K of Fig. 2. Because the signature of K is 16, K is non-amphicheiral. The following fact will be proved in Section 4.

Lemma 2. K is a prime knot.

4. Proof of Lemma 2

We will prove Lemma 2 by the method of Kirby and Lickorish [KL]. A tangle is a set of disjoint two arcs properly embedded in a 3-ball. The tangle as shown in Fig. 3 will be called a trivial tangle. A tangle T in a 3-ball B is prime if it has the following properties.

(i) Any 2-sphere in B , which meets T transversely in two points, bounds in B a ball meeting T in an unknotted spanning arc.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

(ii) The arcs of T cannot be separated by a disc properly embedded in B .

The knot K is constructed by the tangles T_1 and T_2 as shown in Fig. 4. To establish that K is prime, we have only to show that the tangles T_1 and T_2 are prime. If T_i ($i = 1, 2$) did not satisfy property (i) or property (ii), then there would be a ball meeting T_i in a knotted spanning arc as shown in Fig. 5. However, a trivial tangle can be added to T_i on the outside of the ball in Fig. 4 to create the knot K_i ; K_1 is the pretzel knot $K(-2, 3, 7)$, and K_2 is the torus knot of type $(3, 7)$, denoted by $T(3, 7)$. Thus K_i would have $T(2, 5)$ as a factor of its prime decomposition. Let $\Delta_i(t)$ and $\Delta(t)$ be the Alexander polynomials of K_i and $T(2, 5)$ respectively, then $|\Delta_i(-1)| = 1$ must be divisible by $|\Delta(-1)| = 5$. This contradiction establishes Lemma 2.

Remark. K is concordant to $K(-2, 3, 7) \# T(3, 7)$.

5. Question

Using the method of Viro [V], we can construct presumably non-amphicheiral knots; they are prime and the 2-fold covering spaces branched over them are symmetric. The knot of Fig. 6 is one of them. The 2-fold branched covering space is the manifold as shown in Fig. 7. Is this knot non-amphicheiral?




Fig. 6




Fig. 7

References

- [BGM] J. S. Birman, F. González-Acuña and J. M. Montesinos: Heegaard splittings of prime 3-manifold are not unique, *Michigan Math. J.*, 23 (1976), 97-103.
- [K] R. Kirby: Problems in low dimensional manifold theory, *Proc. Symp. Pure Math.* 32, Part 2, 1978, 273-312.
- [KL] R. Kirby and W. B. R. Lickorish: Prime knots and concordance, *Math. Proc. Camb. Phil. Soc.*, 86 (1979), 437-441.
- [M1] J. M. Montesinos: Variedades de Seifert que son recubridores ciclicos ramificados de dos hojas, *Bol. Soc. Mat. Mexicana* (2), 18 (1973), 1-32.
- [M2] _____: Surgery on links and double branched covers of S^3 , *Ann. of Math. Stud.* #84, Princeton Univ. Press, 1975, 227-259.
- [P] R. L. Parris: Pretzel knots, Ph. D. Thesis, Princeton Univ., 1978.
- [R] D. Rolfsen: *Knots and Links*, Publish or Perish Inc., Berkeley, 1976.
- [Sc] H. Schubert: Die eindeutige Zerlegbarkeit eines Knoten in Primknoten, *S.-B. Heidelberger Akad. Wiss. Math.-Natur Kl.*, 3 (1949), 57-104.
- [Se] H. Seifert: Topologie dreidimensionaler gefaseter Räume, *Acta Math.*, 60 (1933), 147-238.
- [Ta] M. Takahashi: Two knots with the same 2-fold branched covering space, *Yokohama Math. J.*, 25 (1977), 91-99.
- [Tr] H. F. Trotter: Some knots spanned by more than one unknotted surface of minimal genus, *Ann. of Math. Stud.* #84, Princeton Univ. Press, 1975, 51-62.
- [V] O. Ya. Viro: Nonprojecting isotopies and knots with homeomorphic coverings, *J. Soviet Math.*, 12 (1979), 86-96 (English translation).
- [W1] F. Waldhausen: Eine Klasse von 3-dimensionalen Mannigfaltigkeiten II, *Invent. Math.*, 4 (1967), 87-117.
- [W2] _____: Über Involution der 3-Sphäre, *Topology*, 8 (1969), 81-91.